

S2 Table. Sensitivity analysis: REACH versus usual care

Variable	REACH (n=19)	Usual care (n=40)	Comparison
Age (median/IQR)	63 (56-65)	60 (52-67)	p 0.603
Gender (n, %)			
• Male	14 (73.7)	24 (60.0)	
• Female	5 (26.3)	16 (40.0)	
ICU LOS (median/IQR)	10 (5-21)	14 (9-22)	p 0.155
Hospital LOS (median/IQR)	23 (14-35)	42 (25-66)	p 0.003*
MV days (median/IQR)	8 (4-14)	13 (5-19)	p 0.245
Admission category (n, %)			
• Acute	17 (89.5)	24 (60.0)	
• Elective	2 (10.5)	16 (40.0)	
Discharge location			
• Home	19 (100)	24 (60.0)	
• Rehabilitation facility	-	16 (40.0)	
SNAQ65+			
• Green	1 (5.3)	3 (7.5)	
• Orange	2 (10.5)	4 (10.0)	
• Red	16 (84.2)	33 (82.5)	

SD: Standard Deviation, LOS: Length Of Stay, IQR: Interquartile Range, MV: Mechanical Ventilation, SNAQ65+:

Short Nutritional Assessment Questionnaire 65+, PImax: maximum static inspiratory mouth pressure, PEmax: maximum static expiratory mouth pressure, GS: Grip Strength, TMST: Two-Minute Step Test

* Significant between group difference, as tested with the independent samples Mann Whitney U test